



CANADIAN WILDLIFE FEDERATION  
FÉDÉRATION CANADIENNE DE LA FAUNE

# AQUACULTURE



CANADIAN WILDLIFE FEDERATION  
FÉDÉRATION CANADIENNE DE LA FAUNE

[CanadianWildlifeFederation.ca](http://CanadianWildlifeFederation.ca)

The purpose of this report is to clearly establish the Canadian Wildlife Federation's position on finfish aquaculture. Moreover, this report demonstrates that our position against open finfish aquaculture is based on thorough research, using the latest and best available knowledge. As such, this report is a living document. As new research is conducted, CWF will update this bibliography and use it to inform our position and messages on the subject.

As you peruse the document, you'll notice that the first section identifies CWF's position and key messages relating to finfish aquaculture and its effects on wildlife. The second section is an annotated bibliography that provides a brief description of scientific research that has investigated the effect of finfish aquaculture on wildlife and provides the reference to the primary scientific literature to allow anyone to examine this research themselves.

I hope you find this report enlightening,

Sean Brilliant  
Manager, Marine Programs  
Canadian Wildlife Federation



# AQUACULTURE

Introduction



THINKSTOCK

# AQUACULTURE

## Introduction

### Q What is Aquaculture?

**A** Aquaculture is the farming of aquatic wildlife like fish, crustaceans, molluscs and aquatic plants. As opposed to commercial fishing, in which fishermen harvest wild fish, aquaculture grows freshwater and saltwater species in controlled conditions.

There are various kinds of aquaculture, but the two types of aquaculture this report will highlight in particular are:

- Closed-containment finfish aquaculture (CCFA)
- Open-pen finfish aquaculture (OPFA)

### Q What is Closed-containment Finfish Aquaculture (CCFA)?

**A** Closed-containment finfish aquaculture (CCFA), is a farming technique that raises fish in a containment system that is a barrier between farms and the natural environment.

There are several challenges for CCFA operations to become a viable means for raising finfish:

- There are additional operational costs and increased fossil fuel requirements
- The species being raised require different (domestication) traits than those currently used
- Influent-effluent treatments need to be developed
- There are substantial capital costs

Although there are some CCFA operations being developed, it is currently viewed as an experimental and developing industry as the challenges listed above are addressed. As it stands, a thorough evaluation of CCFA was done by Chadwick et al in 2010 and the Atlantic Salmon Federation is investing in research to develop CCFA.



### ➔ The Canadian Wildlife Federation's Position on Closed-containment finfish aquaculture

CWF is supportive of research into CCFA technologies and encourages this method of finfish farming to be developed.

# AQUACULTURE

## Introduction



THINKSTOCK

### **Q** What is Open-pen Finfish Aquaculture?

**A** Open-pen finfish aquaculture (OPFA) is the farming of fish species in marine or freshwater within net pens or cages that are open to the natural environment.

### **→** The Canadian Wildlife Federation's Position on Open-pen Finfish Aquaculture

Open-pen finfish aquaculture alters important aquatic habitats and significantly affects native populations of wild salmon and other wildlife. These effects may put the survival of both Pacific and Atlantic salmon species at risk. Although the aquaculture industry creates economic benefits, because of its significant impact on wildlife, CWF would like to see the practice of OPFA in Canada phased out over the next 10 years. In the meantime, CWF believes it imperative that no more OPFA operations be established in Canada.



# AQUACULTURE

## How Open-pen Finfish Aquaculture Affects Wildlife

# AQUACULTURE

## How Open-pen Finfish Aquaculture Affects Wildlife

### → Ecological Interactions



#### *Ecological Interactions*

OPFA operations attract predators such as seals and birds, thereby altering their natural behaviour and putting them at risk of becoming injured in nets or equipment.

#### *Sustainable use of wild resources*

Salmon aquaculture is an added pressure to wild fish stocks. Industry and scientists estimate that it takes between 1.7 kg to 4.9 kg of wild forage fish to produce one kilogram of salmon. While this ratio has significantly dropped in the last few decades, it still remains that more wild fish are required to produce a given amount of salmon.



Furthermore, more than half of the fishmeal produced by global fisheries for forage fish is used in aquaculture. Some species of finfish, like carp and tilapia, don't need nearly as much fish meal and fish oil in their diet as do salmon to produce a similar amount of production.

### → Genetic Consequences

#### *Impacts on Local Salmonids*

Large-scale escapes from OPFA operations have occurred on both the east coast and west coast of Canada. When these salmon escape, there's a risk that they could breed with wild salmon causing reduced genetic diversity and reduced fitness in wild populations. Furthermore, when open-pen farmed salmon escape, wild salmon are faced with a new competitor for resources. Increasingly, research shows this happening to wild salmon as well as the negative effects it's having on their populations.



# AQUACULTURE

## How Open-pen Finfish Aquaculture Affects Wildlife

### → Disease and Parasites

#### *Diseases*

There are several diseases that salmon held in OPFA operations can contract including:

- Bacterial kidney disease. A chronic bacterial disease that is slowly progressive and frequently fatal.
- Infectious haematopoietic necrosis. An infectious viral disease that may infect salmon of any age but predominately young fish. The virus attacks the blood forming tissues of the fish, as well as its kidney and spleen. To date there is no treatment or cure for the disease.
- Infectious salmon anaemia. A viral disease that causes severe anemia in fish.

Fish in OPFA operations can contract diseases from wild fish but, on the other hand, infected fish in OPFA operations can spread diseases to local wild fish populations and can contribute to the spread, intensity and maintenance of these diseases in wild populations.

#### *Parasites*

Sea lice (parasitic copepods that feed on the skin, mucous and blood of fish) can cause morbidity and even mortality in some populations of salmon. Sea lice are natural parasites and wild populations can have large infections un-associated with OPFA. However, fish in OPFA operations can skyrocket the transmission of parasites as they are a year-round host population for the parasites. Even juveniles can be infected in OPFA operations since they remain with the adult population (in the wild, juveniles are usually protected from infection because of their separation from the adult population). Sublethal exposure to sea lice at early life stages reduces the growth of individuals, causing negative effects on populations.



PHOTOS: WIKIMEDIA COMMONS

# AQUACULTURE

## How Open-pen Finfish Aquaculture Affects Wildlife

### → Habitat Alteration

#### *Pharmaceutical pesticides and disinfectants*

Finfish in OPFA operations are treated with therapeutants (various compounds to protect their health). These treatments are usually regulated, however, emergency treatments are required sometimes and those treatments can be difficult to regulate. The potential effects of therapeutants being released into the environment are a concern. They can have terrible effects on other species. Crustaceans, for instance, tend to be particularly sensitive to these treatments – even more so than other species. Research shows that when OPFA therapeutants are used properly, they pose little threat to local populations and crustaceans, however, in some situations; evidence suggests they may have lethal and sublethal effects on local populations.



The use of antibiotics in OPFA operations has also altered local bacterial diversity, increased populations of antibiotic-resistant bacteria, and led to the emergence of new strains of bacteria.

#### *Pollution and habitat alteration*

Fish wastes (faeces, uneaten food, mortalities) accumulate beneath cages of OPFA operations, and can actually modify benthic structure as well as lead to eutrophic (waters rich in nutrients that lead to a boost in plant life, especially algae) and anoxic (low oxygen) conditions. Large-density OPFA operations release quantities of nutrients (nitrogen and phosphorus) comparable to the human sewage production from coastal urban centres.

# AQUACULTURE

## How Open-pen Finfish Aquaculture Affects Wildlife

Moreover, heavy metals with toxic properties, particularly copper, accumulate in the sediments around OPFA operations, often to concentrations above those predicted to cause ecological effects.

OPFA operations provide novel and varied surfaces that organisms may colonize, sometimes in unnaturally large numbers (biofouling). This can lead to other problems, like a boom in invasive species, particularly in the treatment of this issue.

Many of these effects are influenced by the depths, tides and currents of local waters (known as hydrographic conditions). The effects on the community of organisms that live in this environment (benthos) are larger in area and less acute for OPFA operations in deep water compared to those in shallower water.




### *Light*

The lighting used around OPFA operations attracts wild fish to the cages, exacerbating other potential effects, particularly the transmission of disease.



### *Noise*

Operators use noise-makers to deter predators from OPFA cages. While studies show they are ineffective on most target species, like seals and birds, we know they do alter the behaviour and distribution of non-target species, particularly cetaceans (whales, dolphins and porpoises). To date, there is no evidence that these effects on non-target species are long-term or damaging.



# AQUACULTURE

Annotated Bibliography of the Conservation Issues  
of Open-pen Finfish Aquaculture | August 2013

# AQUACULTURE

## Annotated Bibliography of the Conservation Issues of Open-pen Finfish Aquaculture | August 2013

### General Impacts

- Salmon farming is conducted almost exclusively in net pens. A number of environmental impacts can be attributed to this type of aquaculture, as cages are open systems that allow for the release of nutrients, pathogens, and chemical inputs to the marine environment.
- Impacts are commonly grouped into four categories: ecological interactions, genetic consequences, disease and parasites, habitat alteration. More specifically: benthic impacts and alteration, chemical inputs (antibiotics, anti-foulants, pesticides), nutrient loading and deterioration of the benthos, attraction of other organisms and predator exclusion, feed sources, effects of escapees and use of exotic species, and exchange of pathogens (sea lice) (Hutchings et al. 2012).
- An analysis of wild salmon mortality in rivers adjacent to salmon farms demonstrated a large reduction in survival and returns in populations exposed to aquaculture. Mortality increased by more than 50 per cent for each generation for wild salmon exposed to aquaculture (Ford and Myers, 2008).
- On the west coast, only pink salmon showed significant declines correlated with aquaculture (Ford & Myers 2008). This was attributed to the genetic impacts from interbreeding with escaped farm salmon, in addition to disease and other impacts described below.

### Escapees & Genetic Impacts

- Large-scale escapes are frequent occurrences with open-net pen fish farming and can happen through routine handling, or large-scale episodic events such as storms (McGinnity et al. 2003, Naylor et al. 2005), increasing the risk of interactions between wild and farmed salmon. Escapes can have severe impacts on the persistence of wild salmon populations (Morris et al. 2008).
- Morris et al. (2008) compiled a series of studies and found that escaped farmed salmon have been reported in 87 per cent of the watersheds that have been investigated since the inception of the salmon aquaculture industry in 1980 (the majority of the investigated rivers were located in primary aquaculture regions, i.e., N.B. and ME).
- Carr & Whoriskey (2006) assessed the occurrence of escapees from freshwater hatcheries and found escapees in streams every year and escapee smolts outnumbered wild smolts in seven out of the eight years studied in the Magaguadavic River. Escapes of juvenile salmon occurred in streams next to at least 75 per cent of the commercial salmon hatcheries in New Brunswick.

# AQUACULTURE

## Annotated Bibliography of the Conservation Issues of Open-pen Finfish Aquaculture | August 2013

- In addition to increased competition for food and habitat, disease transmission, and overall reduced health, there exists the potential of escaped farm salmon to prey on and breed with wild fish (Flemming et al. 2000). Interactions can also alter the functional response of wild fish to predators (Naylor et al. 2005).
- In contrast to their wild counterparts, farmed salmon are selectively bred for, among other characteristics, faster growth. Due to these genetic differences interbreeding is a major concern.
- Genetic effects of interbreeding between farmed and wild fish may be direct or indirect. Direct genetic effects include the alteration of the wild genome (introgression) as a result of interbreeding between wild and domesticated fish, or the production of sterile hybrids. Indirect effects include the effect of reduced effective population size or altered selection pressure arising from competition (Tymchuk et al. 2006).
- Interbreeding can have genetic effects that threaten the reproductive capability and recovery potential of wild populations, a result of the production of fish that are less fit, as well reduced lifetime success (McGinnity et al. 2003, Thorstad et al. 2008, Hutchings et al. 2012).
- Introgression can eventually lead to genetic homogenization, resulting in an irreversible loss in the genetic variation of wild populations, and subsequently can alter the genetic integrity of native salmon populations, including loss of local adaptation, or the reduced ability to respond to changing environments (Naylor et al. 2005, Fraser et al. 2010, Bourret et al. 2011).
- Farmed Atlantic salmon (*Salmo salar*) tend to be competitively and reproductively inferior to their wild counterparts, with less than one-third the reproductive success (Fleming et al. 2000, Naylor 2005, Hindar et al. 2006). Despite their decreased ability to compete, the farm fishes still were able to compete with the native population. Although breeding performance tends to be inferior to wild salmon, interbreeding does occur successfully resulting in hybridization.
- Hybridization may have contributed to the decline and lack of recovery of many wild Atlantic salmon populations in the Northwest Atlantic because it can change the genetic make-up and life history traits of wild populations (Fraser et al. 2010).
- Hindar et al. (2006) modeled the future of wild salmon populations experiencing invasions of escaped farmed salmon. Results demonstrated that high intrusion could eventually lead to populations that are mixtures of hybrid and farmed descendants within only a few generations and recovery of wild populations in this scenario is not likely.
- Liu et al. (2013) also modeled this showing that increased numbers of escapes reduces the social welfare of the harvest and the wild stocks (recreational).
- McGinnity et al. (2003) demonstrated that interaction of farmed salmon with wild salmon results in lowered fitness, with repeated escapes causing cumulative fitness depression and potentially an extinction vortex in vulnerable populations.

# AQUACULTURE

## Annotated Bibliography of the Conservation Issues of Open-pen Finfish Aquaculture | August 2013

- Most differences appear to result from non-additive gene interactions (rendering effects unpredictable), which affects traits related to fitness (Roberge et al. 2008). These results suggest that interbreeding of fugitive farmed salmon and wild individuals could substantially modify the genetic control of gene transcription in natural populations exposed to high migration from fish farms, resulting in potentially detrimental effects on the survival of these populations (see also Oke et al. 2013).
- The probability that escaped Atlantic salmon will establish populations where the species is exotic seems low, but the possibility cannot be ruled out. Where native populations of salmonids are currently depressed or in decline, conditions for the establishment of Atlantic salmon may be more favourable now than in the past (Thorstad et al. 2008). However, the likelihood of successful hybridization between Atlantic salmon and Pacific salmonid species seems small.
- Feral Atlantic salmon were reported in coastal rivers of B.C.. Volpe et al. (2000) report the first case of successful Atlantic salmon reproduction on the northeast Pacific, which demonstrates the potential for successful colonization, and thus potential for competition for space and food with wild populations of Pacific salmon (in particular *Oncorhynchus spp.*), although these inferences are speculative at best.
- Due to artificial selection of growth in farmed salmon, they tend to have a size advantage over wild salmon and, thus a competitive advantage over wild juveniles (Naylor et al. 2005)
- Stress to wild populations caused by introductions of farmed salmon can affect mortality and growth (Naylor et al. 2005).

### Diseases

- The prevalence of disease is increased with high density of fish in pens commonly observed in salmon aquaculture, and spreads to wild fish, subsequently threatening the persistence of wild fish populations (Hutchings et al. 2012).
- Infectious salmon anaemia (ISA) is one of the most serious infectious diseases facing the economic viability of Atlantic salmon farming (Hammell & Dohoo 2005). It is the most problematic because once infected, there is no treatment (Nylund et al. 1994).
- Characterized by severe anaemia and high mortality, ISA tends to only be diagnosed after fish have been transferred to sea or have been exposed to seawater hatcheries (Nylund et al. 1994).
- While a potential mechanism for spread of disease is passive transmission in seawater, exposure to organic material (i.e., blood) infected with the ISA agent, or to sea lice infected with the ISA virus generated the highest rate of mortality (Nese & Enger 1993, Nylund et al. 1994, Barker et al. 2009). Transfer can occur via skin (mucous membrane, abrasions), urine and feces.

# AQUACULTURE

## Annotated Bibliography of the Conservation Issues of Open-pen Finfish Aquaculture | August 2013

- Miller et al. (2011) discovered genomic signature in Fraser River Sockeye salmon correlated with survival. They hypothesized that the genomic signal associated with elevated mortality is in response to a virus infecting fish before river entry and that persists to the spawning areas (linkages cited with genes associated with leukemia).
- It's been determined that pathogens can be vertically transmitted or transmitted via infected embryos (transmitted from parent to offspring) (Vike et al. 2009).
- IHN disease (infectious haematopoietic necrosis) is an epidemic in farmed Atlantic salmon in B.C. that is endemic in Pacific wild populations. It causes necrosis of the haematopoietic tissues, and ultimate mortality in fish. Farming practices contributed to its spread. However, waterborne transmission may have played a role between farms located in close proximity to one another (Saksida 2006). Similar to ISA, it is spread through feces, urine and mucous.

### Sea lice (parasites)

- The fishes in sea pens can become infected with pathogens like sea lice by wild populations and in turn become point sources for parasites that multiply in crowded pens and spread to wild fish. Floating sea cages (open-net pen) allows free movement of pathogens between wild and farmed finfish (Costello 2009).
- Sea lice are parasitic copepods, ubiquitous on farmed and wild adult salmonids, particularly adult females, in the oceans of the northern hemisphere, that feed on skin, mucous, and blood and are capable of causing host morbidity and mortality (Costello 2009, Saksida et al. 2007a, 2007b).
- *Caligus clemensi* and *Lepeophtheirus salmonis* are two species of sea lice that are commonly found on farmed salmon (Beamish et al. 2009).
- Natural infection of sea lice, consisting mainly of *C. clemensi*, discovered on juvenile Pacific salmon (all species) of the Gulf Islands, BC. They concluded that this infection was not from a fish farm, as the closest was 100 km from study area (Beamish et al. 2009). This confirms the occurrence of natural infection that could be rapidly transmitted in high-density farm cages.
- Migrating adults are separated from juveniles in early life stages, which acts as a barrier to pathogen transmission between age classes (for a time) (Krkošek et al. 2007). The benefit of this is that juveniles are not subject to pathogens until they are older and less vulnerable.
- Winter treatment of farms with parasiticides prior to out-migration of wild juvenile salmon can reduce parasite loads on the wild fish though it was noted that long-term sustainability of this practice may be affected by the ecological effects of the parasiticide and the potential for resistance in the parasites (Peacock et al. 2013, Rogers et al. 2013).

# AQUACULTURE

## Annotated Bibliography of the Conservation Issues of Open-pen Finfish Aquaculture | August 2013

- Intensive open net-pen salmon aquaculture can undermine this natural barrier to transmission by providing a year-round host population in the near-shore marine environment whose pathogens can spill over to sympatric wild juvenile salmonids (Costello 2009). Sub-lethal exposure to lice during the critical early marine period reduces Coho smolt growth with consequences for Coho population dynamics.
- Sea lice transmission from salmon farms may indirectly influence the health of sympatric Coho salmon smolts (Connors et al. 2010) and juvenile pink salmon (Brauner et al. 2013) via the accumulation of lice from the infected prey. Connors et al. (2010) demonstrated that Coho populations had depressed productivity sevenfold when exposed to louse infestations associated with salmon farms, suggesting that parasite transmission from farmed to wild salmon can propagate up a salmonid food web with negative consequences for predatory salmon populations and the ecosystems in which they are embedded. Brauner et al. (2012) showed increased infection affects swimming ability of pink salmon.
- Krkošek et al. (2011a) found that survival was negatively correlated with sea lice abundance on farms in Broughton Archipelago region of B.C. for both pink and Coho salmon. This is a matter up for debate as other studies have reached contradictory conclusions. Marty et al. (2010) did not find a negative correlation between fish farms, or number of sea lice, and pink salmon productivity.
- Wild populations exposed to salmon farms in the Broughton Archipelago, B.C. show a sharp decline in productivity during sea lice infestations relative to pre-infestation years, potentially resulting in declines in pink salmon populations (Krkošek & Hilborn 2011).
- Price et al. (2010) found evidence that salmon farms are a major source of sea lice infestations for wild juvenile pink (*Oncorhynchus gorbuscha*) and chum salmon (*O. keta*) in many aquaculture regions of B.C..
- Krkošek et al. (2011b) examined how sea lice infestation affects predation risk and mortality of juvenile pink and chum salmon and found evidence that lice make juvenile salmon more prone to predation. Infected fish were eaten significantly more often than fish that were not infected, as predators will selectively consume infected prey. Infestation also alters schooling behaviour, as infected fish will occupy peripheral positions in the school. Conversely, predation on infected fish can serve to reduce sea lice abundance in wild fish.
- Krkošek et al. (2005) studied infections of sea lice juvenile pink and chum salmon as they passed an isolated salmon farm during their seaward migration down two long and narrow corridors. Results suggested that the infection pressure imposed by the farm was four orders of magnitude greater than ambient levels, resulting in a maximum infection pressure near the farm that was 73 times greater than ambient levels and exceeded ambient levels for 30

# AQUACULTURE

## Annotated Bibliography of the Conservation Issues of Open-pen Finfish Aquaculture | August 2013

kilometres along the two wild salmon migration corridors. Additionally, sea lice already infecting the wild juveniles were able to reproduce during their migration and re-infect the juveniles increasing the range of impact to 75 km. Amplified sea lice infestations due to salmon farms are a potential limiting factor to wild salmonid conservation with potential impacts on wild salmon fisheries, as well (Liu et al. 2011).

- Krkošek et al. (2013) did a meta-analysis of experimental paraciticide treatments on smolts that, lice infection causes 39 per cent loss of adult salmon recruitment (see also Torrissen et al. 2013).
- Price et al. (2011) demonstrated a potential role of open net-pen salmon farms in transmission of sea lice to wild juvenile sockeye salmon as they migrate from the Fraser River. They found that juveniles were primarily the fish infected with *C. clemensi*. They concluded that salmon farms are elevating levels of sea lice on Fraser River salmon during their critical early migration.
- Other parasites have also been shown to transmit between cultured and wild fish. Mladineo et al. (2013) showed this using gene-flow of a monogenean.

### Antibiotics, Pesticides & Disinfectants

- Although the use of therapeutants is tightly regulated and can only be used if prescribed by a licensed veterinarian, farmers do rely on emergency treatments to combat bacterial infections and ecto-parasites, like sea lice.
- Salmon farmers in Norway, Scotland, Chile and Canada have used a variety of compounds over the years to treat infected salmon, including amoxicillin, florfenicol, tribrisen, oxytetracycline, erythromycin, pyrethrins, hydrogen peroxide, azamethiphos, cypermethrin, ivermectin, emamectin benzoate and teflubenzuron (Burrige et al. 2008, 2010).
- While many of the antibiotics have low toxicity to marine fishes and mammals, crustaceans tend to be more sensitive (McLeese et al. 1980, Davis 1985, Burrige et al. 2004). Therefore, the use of chemicals causes concern among environmentalists and others associated with commercial fisheries. Because anti-lice treatments lack specificity, the effluent released into the marine environment has the potential to cause adverse impacts not only on wild salmon populations, but commercially important species such as the American lobster (*Homarus americanus*) (Johnson et al. 2004, Burrige et al. 2008, 2010).
- Diflubenzuron (crusticide) can be taken up by Atlantic cod but is rapidly cleared and showed to have small effects (Olsvik et al. 2013).
- In general, demonstrated effects of this include alteration of local diversity, increased resistance to antibiotics, and subsequent emergence of new strains of bacteria (Burrige et al. 2010).
- Salmosan® (active ingredient azamethiphos), currently approved for emergency use in N.B., has been shown to have lethal and sublethal effects with repeated exposure on lobster at certain

# AQUACULTURE

## Annotated Bibliography of the Conservation Issues of Open-pen Finfish Aquaculture | August 2013

- concentrations and time of year (BurrIDGE et al. 2008). Organisms were more sensitive to repeated exposure at concentrations of 10 µg/L and during the spring. Field studies have shown that a single treatment with Salmosan® has no negative effect on the survival of non-target organisms except when held within the treatment cage (BurrIDGE 2003).
- Soluble compounds (such as azamethiphos) used for parasite control have potential for widespread dispersion. Other chemicals such as antibiotics (oxytetracycline, Tribissen and Florfenicol) and zinc, present in feed (particle-bound substances), are more likely to settle and accumulate under and close to farm sites if currents are low (DFO 2003).
  - Female lobsters are more sensitive to exposure during spawning and molting season (BurrIDGE et al. 2005).
  - SLICE™ (active ingredient emamectin benzoate) is the most common treatment for sea lice and is a fishfeed premix (BurrIDGE et al. 2004).
  - The fishing industry is concerned about the use of this pesticide and its impact on lobsters and crabs foraging beneath salmon cages (released in uneaten food, fish feces, urine), as crustaceans, particularly lobster, will eat the feed and are known to be particularly sensitive to the pesticide (Waddy et al. 2002, Willis & Ling 2003, Waddy et al. 2007a). Although, acute toxicity is low (BurrIDGE et al. 2004), it can result in premature molting and loss of eggs by female American lobsters (Waddy et al. 2002). However, the question remains as to whether or not lobsters are likely to ingest enough of the chemical, based on industry concentrations, to cause any observable effect. Feeding studies have confirmed that lobsters ingest less than the amount required to initiate premature molting (Waddy et al. 2007a, 2007b).
  - In addition, based on the concentration required to kill a lobster, and the small amount of medicated food consumed, researchers have predicted that chronic exposure is unlikely to result in the death of exposed lobsters (BurrIDGE et al. 2004).
  - Alpha Max® (active ingredient Deltamethrin, a pyrethroid insecticide) was approved for emergency use in 2009 and 2010, but is currently not in use. Fairchild et al. (2010) report that “Pyrethroid insecticides are among the most toxic insecticides known” (causing immobility or mortality), and that “among the pyrethroid insecticides, deltamethrin is often the most toxic to crustaceans” (p. iv). In lab conditions, Fairchild et al. (2010) found lobster larvae to be susceptible to deltamethrin at much lower concentration than the recommended treatment dose.
  - Ernst et al. (2001) demonstrated the dispersal capacity of cypermethrin, another widely known highly toxic (particularly for crustaceans) pesticide. Water samples collected from the plume were toxic in a 48 hour lethality test to *Eohaustorius estuarius* for cypermethrin up to five hours. They concluded that single treatments have the potential to affect non-target invertebrates near cage sites.

# AQUACULTURE

## Annotated Bibliography of the Conservation Issues of Open-pen Finfish Aquaculture | August 2013

- Alternatively, because pyrethroids tend to adsorb onto particulate matter, chronic exposures may not occur other than in laboratory studies. Cypermethrin absorbed by sediment was not acutely toxic to grass shrimp until concentrations in sediment were increased to the point where partitioning into the overlying water resulted in acutely lethal concentrations (Clark et al. 1987).
- The safety of human food can also directly be affected by the presence of residual antibiotics in farmed fish which have been dosed with antibiotics, or from antibiotics that have leached into the surrounding environment and reached wild fish and shellfish collected for consumption (Samuelson et al. 1992, Coyne et al. 1997, Grave et al. 1999, Cabello 2003, Fortt et al. 2007, White & McDermott 2009).

### Pollution

- Environmental concerns with cage aquaculture are primarily associated with the release of waste materials into the aquatic environment (Rooney & Podemski 2009).
- Fish waste from cages flows into ocean, adding potentially harmful excess nutrients in the ecosystem; uneaten food and faeces can accumulate on the seabed underneath the pens and degrades, oxygen then becomes depleted, all with significant implications for water quality and species composition (eutrophication, release of noxious gases, etc.) (Dean et al. 2007, Hutchings et al. 2012).
- With an annual production of 35,000 tonnes of salmon, approximately 7,200 tonnes of untreated fecal waste, 1,000 tonnes of nitrogen, and 115 tonnes of phosphorus are discharged into coastal waters. In terms of human waste, this equates to an annual equivalent of 93,450 people; the nitrogen discharge is equivalent to that of 437,500 people; and the phosphorus discharge is equal to that of 63,000 people. No other industry is allowed to discharge untreated waste like this into water bodies. Nitrogen and phosphorus are both nutrients that cause algal blooms, which lead to eutrophication, a condition to which very few species are tolerant. Aquaculture has been demonstrated to be by far the largest source of nutrients to the L'Etang - Bliss Harbour – Black's Harbour area of the Bay. It has been proven that a significant number of species have been lost in this area as a result (DFO 2003, Harvey & Milewski 2007).
- Metals enter the marine environment from aquaculture activity either from anti-foulant paints or as constituents of fish food. The diet of farmed Atlantic salmon contains trace metals such as copper, zinc, iron, cadmium, manganese, and others (Dean et al. 2007, Burrige et al. 2010).

# AQUACULTURE

## Annotated Bibliography of the Conservation Issues of Open-pen Finfish Aquaculture | August 2013

- Evaluation for toxicity of sediments under salmon cages in the Bay of Fundy, as well as in Scotland, and at various distances away from the cages demonstrate elevated levels of copper (above the threshold effects level), elevated zinc, other metals, ammonia nitrogen, sulfide, total organic carbon, and other organic compounds (Burridge et al. 1999, Dean et al. 2007, Hargrave 2010).
- In the sediments, highest copper concentrations (32–42 mg Cu/kg) are found consistently at the farm site, moderately elevated concentrations at 400 to 2000 metres distance and the lowest concentrations (6–7 mg Cu/kg) are found at a greater distance from the farm site, suggesting that it is plausible for copper, released for purposes of aquaculture and settled in the sediments, to be transferred upward through the water column to the sea surface microlayer. These concentrations are enriched, and can then travel horizontally at relatively high drift speeds due to winds, leading to footprints with scale of one km or more, with the potential for impacts to persist through years of fallow. Observed concentrations of copper in the sediment exceeded optimal threshold levels for the protection of marine life (Loucks et al. 2012).
- Toxicity tests on several species of marine copepods demonstrated reduced survival and reproduction when exposed to metals, such as silver, zinc, and copper in their algal diets and water (Bielmyer et al. 2006, Lauer & Bianchini 2010).
- Copper exposure reduced egg production and hatching rate in male and female copepods (*Acartia tonsa*). Impacts varied with type of exposure with waterborne eliciting a greater adverse effect on egg production than diet borne (Lauer & Bianchini 2010).
- Sediments enriched in these metals have been linked to decreased production in the clam *Macoma balthica*, due to failed gamete production (Burridge et al. 1999).
- Exposure to copper was linked to chromosomal anomalies in spiny lobster, which increased with exposure to copper, demonstrating adverse impacts at the cellular level, internal organs changed colour, and observed changes in wet weight of muscle, hepatopancreas, and gills (Maharajan et al. 2011). In addition, impacts to the heart were observed, specifically noted was inflammation and cell and tissue damage to inner organs (Maharajan et al. 2012). This could affect vital physiological functions, such as absorption, storage and secretion of the hepatopancreas, digestion of gut and respiration, osmotic and ionic regulations of the gills, which in turn could ultimately affect the survival and growth of *P. homarus* (Maharajan et al. 2012).
- Accumulation of fish feces and waste feed enhances aerobic and anaerobic microbial activity, both representing nitrogen-rich organic matter (Giles 2008), as it stimulates phytoplankton production and increase oxygen demand (Hargrave 2003). Eutrophication can cause shifts in phytoplankton species assemblages, although it has been hard to directly link finfish aquaculture to harmful algal blooms (HABS), alters benthic faunal communities, and reduces fish health, which increases the likelihood of parasitism and disease (Hargrave 2003).

# AQUACULTURE

## Annotated Bibliography of the Conservation Issues of Open-pen Finfish Aquaculture | August 2013

- The degree of impact is dependent on local hydrographic conditions, cage depth, uptake by phytoplankton and other organisms, and re-suspension of material (Hargrave 2003, Borja et al. 2009).
- Intermittent hypoxic events from changes in dissolved oxygen (DO) are occurring in cages on the south coast of Newfoundland. These events can be observed for as long as two and a half months and are mostly happening during the summer season. Fish swimming data show that these events do not cause avoidance behavior, and thus fish are negatively impacted as changes in DO can impact fish feeding and growth (Burt et al. 2012).
- Benthic impact from fish farms is a function of site and farm characteristics. Impacts were less intense but further spread in deep areas, whereas impacts in shallow areas appeared more intense but confined to a smaller area around the farm (Giles 2008).
- A study examining the effects of a rainbow trout farm demonstrated a reduction in benthic invertebrate abundance and species richness below the cage due to high levels of organic loading (Rooney & Podemski 2009). However, the impacts dissipated with distance from cage.
- There is potential for far-field effects of waste and nutrient input from salmon farms. Three types of broad-scale changes distant from farm sites include eutrophication, sedimentation and effects on the food web (Hargrave 2003).
- In addition to the creation of a water surface film from the dust from feed dispensers, uneaten feed and feces contribute to the suspended particulate matter. Potential exists for widespread dispersion and horizontal transport of particulate matter depending on tidal flow, circulation, wind and wave energy, and other physical processes (Hargrave 2003). This can lead to eutrophication events, and in terms of the release of antibiotics and pesticides, can result in inlet-wide scale impacts.
- The presence of such large and varied surfaces that accompanies the aquaculture sector provides for a broad diversity of epibiotic organisms including barnacles, bivalves, bryozoans, polychaetes, ascidians, hydroids, sponges and algae to settle and grow. These marine algae and animals, collectively termed biofouling, are severely problematic to culture operations and can have significant economic impacts (Fitridge et al 2012).
- Fitridge et al. (2012) reported three main negative effects of biofouling:
  - » Restriction of water flow and exchange due to growth of fouling organisms on cages, which reduces water quality and lowers DO
  - » Disease risk due to fouling organisms acting as reservoirs for pathogens, an impact that is exacerbated by stress induced by reduced water flow that leads to reduced immunity to pathogens
  - » Cage deformation and structural fatigue due to weight of fouling organisms, increasing the potential for escapees

# AQUACULTURE

## Annotated Bibliography of the Conservation Issues of Open-pen Finfish Aquaculture | August 2013

### Local Predators

- Due to the high abundance of fish in a confined area, sea cages can attract predators, mainly seabirds and marine mammals (Cottee et al. 2009, Jimenez et al. 2013). In addition to increased mortality, stress, and injury to the farmed fish, this can result in predation of other fish species, as well as economic damages incurred by the farmers.
- Predatory animals like seals, sea lions, sharks, birds, and other marine wildlife can become entangled in the fish pens, and deterrents (AHDs) can alter behaviour of predators (Hutchings et al. 2012).

### Light

- Open net-pen salmon farms in British Columbia routinely illuminate their net-pens during the winter and spring, which has been demonstrated to increase the abundance of wild fish larvae, juveniles, and adults around pens, thereby increasing the probability that farmed fish and wild species directly and indirectly interact in coastal marine environments (McConnell et al. 2010).

### Noise

- AHDs are generally ineffective at deterring predators, specifically seals, over the long-term, as field observations indicate that seals appear to habituate to AHDs, possibly due to hearing impairment (not yet assessed) (DFO 2010).
- Cetaceans such as harbour porpoises and killer whales appear to be sensitive to AHDs and field studies have shown they are displaced from large areas when AHDs are in use. Yet impacts seem to be short-term and localized (e.g., avoidance, masking communication and echolocation sounds), and do not appear sufficient to cause injury to, or permanent displacement of, aquatic animals (DFO 2010).

### Fish as Food

- The finfish aquaculture sector is highly dependent on forage fish for dietary input. Forage fish are reduced to fishmeal and fish oil, which threatens the future of wild populations and can impact other wildlife that depends on this food source (Alder et al. 2008, Tacon & Metian 2008, Hutchings et al. 2012). More than 50 per cent of fishmeal produced is used in the aquaculture sector, China being the major fishmeal consumer. China currently consumes more than 1.3 million tons of fishmeal per year.

# AQUACULTURE

## Annotated Bibliography of the Conservation Issues of Open-pen Finfish Aquaculture | August 2013

- The ratio of wild fisheries inputs to farmed fish outputs for Atlantic salmon remains higher than other farmed species. High amounts of forage fish are used in fishmeal and fish oil to sustain this farmed fishery (Naylor et al. 2009).
- Tacon & Metian (2008) reported that, in 2006, the aquaculture sector consumed 3724 thousand tons of fish meal (68.2 per cent of the total global fish meal production in 2006) and 835 thousand tons of fish oil (88.5 per cent of the total reported fish oil production in 2006), or the equivalent of 16.6 million tonnes of small pelagic forage fish (using a wet fish to fish meal processing yield of 22.5 per cent and wet fish to fish oil processing yield of five per cent).
- It takes 4.9 kg of wild fish to produce one kg of salmon (Tacon & Metian 2008), meaning that producing salmon results in a net loss of fish protein for other species that rely on this resource, including human consumption (Harvey & Milewski 2007).
- The International Fishmeal and Fish Oil Organization (IFFO) presented estimates for Fish In – Fish Out (FIFO) ratios for salmon (Jackson 2009) as 2.2 kg wild fish to one kg of salmon. This was further refined to include global pig and poultry production (because they also use fishmeal), this reduced the FIFO ratio to 1.68 kg wild fish to one kg of salmon.
- Waagbo et al. (2013) fed Atlantic salmon diets with increasing amounts of vegetable oil and showed they maintained health and product quality, though none of the groups showed net fish production (FIFO  $\geq$  1.65).

### Closed-containment Finfish Aquaculture

- Open-sea net pens have far greater potential and realized negative consequences to marine biodiversity than closed-containment facilities (Hutchings et al. 2012).
- The development of closed-containment systems, a term widely used to describe a range of production systems that employ an impermeable barrier to isolate the culture environment from surrounding ecosystems, has been proposed as a viable alternative to open-pen systems (Ayer & Tyedmers 2009).
- The benefits of closed-containment have been reported as follows (Ayer & Tyedmers 2009):
  - » Fish farmers can exert greater control over rearing systems (which, in turn, can improve the quality of fish while minimizing the environmental impact)
  - » Minimized escapes
  - » Minimized predator interactions
  - » Reduced disease transmission
  - » Lower feed inputs
  - » Higher stocking densities
  - » Improved waste management capabilities

# AQUACULTURE

## Annotated Bibliography of the Conservation Issues of Open-pen Finfish Aquaculture | August 2013

- Ayer & Tyedmers (2009) determined that while land-based closed containment systems provide a more environmentally-friendly alternative to the issues associated with open-pen aquaculture (release of fish waste and other nutrients and chemicals into the marine environment), there are tradeoffs to be considered (it is the poorest environmental performance in this study). This system increased contributions to several other environmental impacts of global concern, including global warming, acidification, and abiotic resource use (non-renewable resources), as it uses more fossil fuels to power electricity driven pumps. A floating bag system demonstrated the highest environmental performance relative to the others due to lower feed input per ton, lower energy demand, and a reliance on hydro-power because it is marine-based. However, performance was related to site location.
- Forster & Slaski (2010) determined that there were no closed-containment systems found globally that is producing exclusively adult Atlantic salmon, and many previous attempts to do this failed. The primary cited reasons include mechanical breakdown, poor fish performance, management failure, declines in market price and inadequate financing.
- Operating and capital cost are among the major factors that prevent viable application of this technology to salmon farming (Chadwick et al. 2010).
- Stechey & Robertson (2010) described and examined five types of aquaculture production systems (including conventional net pen, floating closed-confinement systems both with rigid walls and flexible walls, land-based flow-through systems, and land-based recirculation systems).
- They determined that land-based intensive recirculation facilities demonstrated the highest level of containment; the risk of release of pathogens and parasites was categorized as moderate for all systems; and the risk of escapes was characterized as “low” or “negligible” for all systems except the conventional net pen and floating flexible wall system (Stechey & Robertson 2010).
- Land-based, solid-wall, recirculation and reuse technologies (Recirculating Aquaculture Systems) showed promise for rearing of salmon in fresh or brackish water due to the potential for reduced energy costs compared with those associated with the pumping of seawater (Stechey & Robertson 2010).
- In a closed systems there is the potential for increased exposure and the spread of pathogens because fish stocking densities would be higher and water turnover rates lower leading to increased accumulation of waste debris (including pathogens) (Byrne et al. 2010). However, closed systems improve the separation of farmed from wild fish, lowering the risk of spread to wild populations.
- Complete disruption of pathogen exchange between a closed pen facility and the environment is likely only possible in a land-based recirculation system (Byrne et al. 2010).
- Pathogen removal involves the treatment of influent and effluent water flows, the effectiveness of which has yet to be determined (Byrne et al. 2010).

# AQUACULTURE

## References

### References

- Alder J, Campbell B, Karpouzi V, Kaschner K, Pauly D (2008) Forage fish: from ecosystems to markets. *Annu Rev Env Resour* 33: 153-166.
- Ayer NW & Tyedmers PH (2009) Assessing alternative aquaculture technologies: Life cycle assessment of salmonid culture systems in Canada. *J Clean Prod* 17(3): 362-373.
- Barker DE, Braden LM, Coombs MP, Boyce B (2009) Preliminary studies on the isolation of bacteria from sea lice, *Lepeophtheirus salmonis*, infecting farmed salmon in British Columbia, Canada. *Parasitol Res* 105(4): 1173-1177.
- Beamish R, Gordon E, Jones S, Neville C, Lange K, Sweeting R, Wade J, Pennell W (2009) A large, natural infection of sea lice on juvenile Pacific salmon in the Gulf Islands area of British Columbia, Canada. *Aquaculture* 297(1-4): 31-37.
- Bielmyer GK, Grosell M, Brixti KV (2006) Toxicity of silver, zinc, copper, and nickel to the copepod *Acartia tonsa* exposed via a phytoplankton diet. *Environ Sci Technol* 40(6): 2063-8.
- Borja A, Rodriguez JG, Muxika I, Black K, Nickell TD, Bodoy A, Emblow C, Fernandes TF, Forte J, Karakassis I, Papageorgiou N, Sevastou K, Pranovi F, Tomassetti P, Angel D (2009) Assessing the suitability of a range of benthic indices in the evaluation of environmental impact of fin and shellfish aquaculture located in sites across Europe. *Aquaculture* 293(3-4): 231-240.
- Bourret V, Bernatchez L, O'Reilly PT, Carr JW, Berg PR (2011) Temporal change in genetic integrity suggests loss of local adaptation in a wild Atlantic salmon (*Salmo salar*) population following introgression by farmed escapees. *Heredity* 106(3): 500-510.
- Brauner CJ, Sackville M, Gallagher Z, Tang S, Nendick L, Farrell AP (2012) Physiological consequences of the salmon louse (*Lepeophtheirus salmonis*) on juvenile pink salmon (*Oncorhynchus gorbuscha*): implications for wild salmon ecology and management, and for salmon aquaculture. *Philosophical Transactions of the Royal Society B-Biological Sciences* 367:1770-1779
- Burrige LE, Doe K, Haya K, Jackman PM, Lindsay G, Zitko V (1999) Chemical analysis and toxicity tests on sediments under salmon net pens in the Bay of Fundy. *Can Tech Rep Fish Aquat Sci* 2291: 1-39.
- Burrige LE (2003) Chemical use in marine finfish aquaculture in Canada: a review of current practices and possible environmental effects. *Can Tech Rep Fish Aquat Sci* 2450: 1-131.

# AQUACULTURE

## References

- Burridge LE, Hamilton N, Waddy SL, Haya K, Mercer SM, Greenhalgh R, Tauber R, Radecki SV, Crouch LS, Wislocki PG, Endris RG (2004) Acute toxicity of emamectin benzoate (SLICE™) in fish feed to American lobster, *Homarus americanus*. *Aquacult Res* 35(8): 713-722.
- Burridge LE, Haya K, Waddy SL (2005) Seasonal lethality of the organophosphate pesticide, azamethiphos to female American lobster (*Homarus americanus*). *Ecotox Environ Safe* 60(3): 277-81.
- Burridge LE, Haya K, Waddy SL (2008) The effect of repeated exposure to azamethiphos on survival and spawning in the American lobster (*Homarus americanus*). *Ecotox Environ Safe* 69(3): 411-415.
- Burridge L, Weis JS, Cabello F, Pizarro J, Bostick K (2010) Chemical use in salmon aquaculture: A review of current practices and possible environmental effects. *Aquaculture* 306(1-4): 7-23.
- Burt K, Hamoutene D, Mabrouk G, Lang C, Puestow T (2012) Environmental conditions and occurrence of hypoxia within production cages of Atlantic salmon on the south coast of Newfoundland. *Aquacult Res* 43(4): 607-620.
- Byrne P, Westcott J, McClure C, Caraguel C, Hammell L (2010) Mitigating pathogen transmission for commercial solid wall marine salmon farms in British Columbia – a concept paper: disinfection technologies, biosecurity, and animal health. In: Chadwick EMP, Parsons GJ, Sayavong B (eds) *Evaluation of closed-containment technologies for saltwater salmon aquaculture*. NRC Research Press, Ottawa, p 111-130.
- Cabello FC (2003) Antibiotics and aquaculture: an analysis of their potential impact upon the environment, human and animal health in Chile. *Fundacion Terram Analisis de Politicas Publicas* 17: 1-16. Accessed 10 Aug. [www.terram.cl/index](http://www.terram.cl/index).
- Carr JW & Whoriskey FG (2006) The escape of juvenile farmed Atlantic salmon from hatcheries into freshwater streams in New Brunswick, Canada. *ICES J Mar Sci* 63(7): 1263-1268.
- Chadwick EMP, Parsons GJ, Sayavong B (eds) (2010) *Evaluation of closed-containment technologies for saltwater salmon aquaculture*, Vol. NRC Research Press, Ottawa, Ontario, Canada. 160 pp.
- Clark, J.R., Patrick Jr., J.M., Moore, J.C., Loes, E.M., 1987. Waterborne and sediment-source toxicities of six organic chemicals to grass shrimp (*Palaemonetes pugio*) and amphioxus (*Branchiostoma caribaeum*). *Arch Environ Con Tox* 16(4): 401-407.
- Connors B, Krkosek M, Ford J, Dill L (2010) Coho salmon productivity in relation to salmon lice from infected prey and salmon farms. *J Appl Ecol* 47(6): 1372-1377.

# AQUACULTURE

## References

- Costello MJ (2009) How sea lice from salmon farms may cause wild salmonid declines in Europe and North America and be a threat to fishes elsewhere. *P Roy Soc B-Biol Sci* 276(1672): 3385-3394.
- Cottee S & Petersan P (2009) Animal welfare and organic aquaculture in open systems. *J Agr Environ Ethics* 22(5): 437-461.
- Coyne R, Hiney M, Smith P (1997) Transient presence of oxytetracycline in blue mussels (*Mytilus edulis*) following its therapeutic use at a marine Atlantic salmon farm. *Aquaculture* 149: 175-181.
- Davis JH (1985) The pyrethroids: an historical introduction. In: Leahey JP (ed) *The pyrethroid insecticides*. Taylor and Francis Ltd., London, 1-41.
- Dean RJ, Shimmield TM, Black KD (2007) Copper, zinc and cadmium in marine cage fish farm sediments: an extensive survey. *Environ Pollut* 145(1): 84-95.
- DFO (Department of Fisheries and Oceans) (2003) Salmon holding capacity in Southwestern New Brunswick. *Can Tech Rep Fish Aquat Sci* 2489: 1-60. Accessed 9 Aug. <http://www.dfo-mpo.gc.ca/Library/275378.pdf>.
- DFO (Fisheries and Oceans Canada) (2010) Pathways of effects for finfish and shellfish aquaculture. *DFO Can Sci Advis Sec Sci Advis Rep* 2009/071: 1-22. Accessed 9 Aug. [http://www.dfo-mpo.gc.ca/CSAS/Csas/Publications/SAR-AS/2009/2009\\_071\\_e.pdf](http://www.dfo-mpo.gc.ca/CSAS/Csas/Publications/SAR-AS/2009/2009_071_e.pdf)
- Ernst W, Jackman P, Doe K, Page F, Julien G, Mackay K, Sutherland T (2001) Dispersion and toxicity to non-target aquatic organisms of pesticides used to treat sea lice on salmon in net pen enclosures. *Mar Pollut Bull* 42: 433-444.
- Fairchild WL, Doe KG, Jackman PM, Arsenault JT, Aubé JG, Losier M, Cook AM (2010) Acute and chronic toxicity of two formulations of the pyrethroid pesticide deltamethrin to an amphipod, sand shrimp and lobster larvae. *Can Tech Rep Fish Aquat Sci* 2876: 1-34.
- Fitridge I, Dempster T, Guenther J, de Nys R (2012) The impact and control of biofouling in marine aquaculture: a review. *Biofouling* 28(7): 649-669.
- Fleming I, Hindar K, Mjølnerød I, Jonsson B, Balstad T, Lamberg A (2000) Lifetime success and interactions of farm salmon invading a native population. *Proc Biol Sci* 267(1452): 1517-1523
- Ford JS & Myers RA (2008) A global assessment of salmon aquaculture impacts on wild salmonids. *PloS Biol* 6(2): 0411-0417.

# AQUACULTURE

## References

- Forster J & Slaski R (2010) Review of past experience. In: Chadwick EMP, Parsons GJ, Sayavong B (eds) Evaluation of closed-containment technologies for saltwater salmon aquaculture. NRC Research Press, Ottawa, p 5-25.
- Fortt ZA, Cabello FC, Buschmann RA (2007) Residues of tetracycline and quinolones in wild fish living around a salmon aquaculture center in Chile. *Rev Chil Infectol* 24: 14–18.
- Fraser DJ, Houde ALS, Debes PV, Hutchings JA, O'Reilly P, Eddington JD (2010) Consequences of farmed-wild hybridization across divergent wild populations and multiple traits in salmon. *Ecol Appl* 20(4): 935-953.
- Giles H (2008) Using Bayesian networks to examine consistent trends in fish farm benthic impact studies. *Aquaculture* 274(2-4): 181-195
- Grave K, Lingaas E, Bangen M, Ronning RM (1999) Surveillance of the overall consumption of antibacterial drugs in humans, domestic animals and farmed fish in Norway in 1992 and 1996. *J Antimicrob Chemoth* 43: 243-252.
- Hammell KL & Dohoo IR (2005) Mortality patterns in infectious salmon anaemia virus outbreaks in New Brunswick, Canada. *J Fish Dis* 28(11): 639-50.
- Hargrave BT (2010) Empirical relationships describing benthic impacts of salmon aquaculture. *Aquacult Environ Interact* 1: 33-46.
- Hargrave BT (2003) Far-field environmental effects of marine finfish aquaculture. *Can Tech Rep Fish Aquat Sci* 2450: 1-49.
- Harvey J & Milewski I (2007) Salmon aquaculture in the Bay of Fundy: an unsustainable industry. Conservation Council of New Brunswick Inc., Fredericton, 1-65.
- Hindar K, Fleming IA, Mcginnity P, Diserud O (2006) Genetic and ecological effects of salmon farming on wild salmon: modelling from experimental results. *ICES J of Mar Sci* 63(7): 1234-1247.
- Hutchings JA, Côté IM, Dodson JJ, Fleming IA, Jennings S, Mantua NJ, Peterman RM, Riddell BE, Weaver AJ, VanderZwaag DL (2012) Sustaining Canadian marine biodiversity: responding to the challenges posed by climate change, fisheries, and aquaculture. Expert panel report prepared for the Royal Society of Canada, Ottawa, p1-315.
- Jackon, A (2009) Fish in–fish out ratios explained. *Aquaculture Europe* 34(3): 5-10.
- Johnson SC, Treasurer JW, Bravo S, Nagasawa K, Kabata Z (2004) A review of the impact copepods on marine aquaculture. *Zool Stud* 43: 229–243.

# AQUACULTURE

## References

- Jimenez JE, Arriagada AM, Fonturbel FE, Camus PA, Avila-Thieme MI (2013) Effects of exotic fish farms on bird communities in lake and marine ecosystems. *Naturwissenschaften* 100:779-787
- Krkošek M, Lewis MA, Volpe JP (2005) Transmission dynamics of parasitic sea lice from farm to wild salmon. *Proc Biol Sci* 272(1564): 689-696.
- Krkošek M, Gottesfeld A, Proctor B, Rolston D, Carr-Harris C, Lewis MA (2007) Effects of host migration, diversity, and aquaculture on sea lice threats to Pacific salmon populations. *P Roy Soc Lond B Bio* 274:3141–3149.
- Krkošek M, Connors BM, Morton A, Dill LM, Lewis MA, Hilborn R (2011a) Effects of parasites from salmon farms on productivity of wild salmon. *P Natl Acad Sci USA* 108(35): 14700-14704.
- Krkošek M, Lewis MA, Hilborn R, Connors BM, Mages P, Dill LM, Ford H, Peacock S, Volpe JP, Ford JS, Alexandra M (2011b) Fish farms, parasites, and predators: implications for salmon population dynamics. *Ecol Appl* 21(3): 897-914.
- Krkošek M & Hilborn R (2011) Sea lice (*Lepeophtheirus salmonis*) infestations and the productivity of pink salmon (*Oncorhynchus gorbuscha*) in the Broughton Archipelago, British Columbia, Canada. *Can J Fish Aquat Sci* 68: 17-29.
- Krkošek M, Revie CW, Gargan PG, Skilbrei OT, Finstad B, Todd CD (2013) Impact of parasites on salmon recruitment in the Northeast Atlantic Ocean. *Proc R Soc B* 280: 20122359. <http://dx.doi.org/10.1098/rspb.2012.2359>
- Lauer MM & Bianchini A (2010) Chronic copper toxicity in the estuarine copepod *Acartia tonsa* at different salinities. *Environ Toxicol Chem* 29(10): 2297-2303.
- Liu Y, Sumaila U, Volpe J (2011) Potential ecological and economic impacts of sea lice from farmed salmon on wild salmon fisheries. *Ecol Econ* 70(10): 1746-1755.
- Liu Y, Diserud OH, Hindar K, Skonhøft A (2013) An ecological-economic model on the effects of interactions between escaped farmed and wild salmon (*Salmo salar*). *Fish and Fisheries* 14:158-173
- Loucks RH, Smith RE, Fisher CV, Fisher EB (2012) Copper in the sediment and sea surface microlayer near a fallowed, open-net fish farm (Article in Press). *Mar Pollut Bull*. Accessed 27 Jul. <http://dx.doi.org/10.1016/j.marpolbul.2012.05.025>.
- Maharajan A, Kumarasamy P, Vaseeharan B, Rajalakshmi S, Vijayakumaran M, Chen JC (2011) Effect of copper on morphology, weight, and chromosomal aberrations in the spiny lobster, *Panulirus homarus* (Linnaeus, 1758). *Biol Trace Elem Res* 144: 769-780.

# AQUACULTURE

## References

- Maharajan A, Rajalakshmi S, Vijayakumaran M, Kumarasamy P (2012) Sublethal effect of copper toxicity against histopathological changes in the spiny lobster, *Panulirus homarus* (Linnaeus, 1758). *Biol Trace Elem Res* 145(2): 201-210.
- Marty GD, Saksida SM, Quinn II TJ (2010) Relationship of farm salmon, sea lice, and wild salmon populations. *P Natl Acad Sci USA* 107(52): 22599-22604.
- McConnell A, Routledge R, Connors BM (2010) Effect of artificial light on marine invertebrate and fish abundance in an area of salmon farming. *Mar Ecol Prog Ser* 419: 147-156.
- McGinnity P, Prodöhl P, Ferguson A, Hynes R, Maoiléidigh NÓ, Baker N, Cotter D, O'Hea B, Cooke D, Rogan G, Taggart J, Cross T (2003) Fitness reduction and potential extinction of wild populations of Atlantic salmon, *Salmo salar*, as a result of interactions with escaped farm salmon. *Proc Biol Sci* 270(1532): 2443-2450.
- McLeese DW, Mecalfe CD, Zitko V (1980) Lethality of permethrin, cypermethrin and fenvalerate to salmon, lobster and shrimp. *B Environ Contam Tox* 25: 950-955.
- Miller KM, Li S, Kaukinen KH, Ginther N, Hinch SG, Hruska KA, Hammil E, Curtis JMR, Patterson DA, Sierocinski T, Donnison L, Pavlidis P, Cooke SJ, English KK, Farrell AP (2011) Genomic signatures predict migration and spawning failure in wild Canadian salmon. *Science* 331(6014): 214-217.
- Mladineo I, Segvic-Bubic T, Stanic R, Desdevises Y (2013) Morphological Plasticity and Phylogeny in a Monogenean Parasite Transferring between Wild and Reared Fish Populations. *Plos One* 8 (4): e62011. doi:10.1371/journal.pone.0062011
- Morris M, Fraser D, Heggelin A, Whoriskey F, Carr J, O'Neil S, Hutchings J (2008) Prevalence and recurrence of escaped farmed Atlantic salmon (*Salmo salar*) in eastern north American rivers. *Can J Fish Aquat Sci* 65: 2807-2826.
- Naylor R, Hindar K, Fleming IA, Goldberg R, Williams S, Volpe J, Whoriskey F, Eagle J, Kelso D, Mangel M (2005) Fugitive salmon: Assessing the risks of escaped fish from net-pen aquaculture. *Bioscience* 55(5): 427-437.
- Naylor RL, Chiu A, Hardy RW, Bureau DP, Hua K, Elliott M, Farrell AP, Forster I, Gatlin DM, Goldberg RJ, Nichols PD (2009) Feeding aquaculture in an era of finite resources. *P Natl Acad Sci USA* 106(36): 15103-15110.
- Nese L & Enger O (1993) Isolation of *Aeromonas salmonicida* from salmon lice *Lepeophtheirus salmonis* and marine plankton. *Dis Aquat Org* 16(1): 79-81.

# AQUACULTURE

## References

- Nylund A (1994) Mechanisms for transmission of infectious salmon anaemia (ISA). *Dis Aquat Org* 19: 95-100.
- Oke KB, Westley PAH, Moreau DTR, Fleming IA (2013) Hybridization between genetically modified Atlantic salmon and wild brown trout reveals novel ecological interactions. *Proceedings of the Royal Society B-Biological Sciences* 280 20131047. <http://dx.doi.org/10.1098/rspb.2013.1047>
- Olsvik PA, Samuelsen OB, Erdal A, Holmelid B, Lunestad BT (2013) Toxicological assessment of the anti-salmon lice drug diflubenzuron on Atlantic cod *Gadus morhua*. *Diseases of Aquatic Organisms* 105:27-43
- Peacock SJ, Krkosek M, Proboszcz S, Orr C, Lewis MA (2013) Cessation of a salmon decline with control of parasites. *Ecological Applications* 23:606-620
- Price MH, Proboszcz SL, Routledge RD, Gottesfeld AS, Orr C, Reynolds JD (2011) Sea louse infection of juvenile sockeye salmon in relation to marine salmon farms on Canada's west coast. *PloS One* 6(2): 1-9. Accessed 27 Jul. doi:10.1371/journal.pone.0016851.
- Price M, Morton A, Reynolds J (2010) Evidence of farm-induced parasite infestations on wild juvenile salmon in multiple regions of coastal British Columbia, Canada. *Can J Fish Aquat Sci* 67: 1925-1932.
- Roberge C, Normandeau E, Einum S, Guderley H, Bernatchez L (2008) Genetic consequences of interbreeding between farmed and wild Atlantic salmon: insights from the transcriptome. *Mol Ecol* 17(1): 314-324.
- Rogers LA, Peacock SJ, McKenzie P, DeDominicis S, Jones SRM, Chandler P, Foreman MGG, Revie CW, Krkosek M (2013) Modeling Parasite Dynamics on Farmed Salmon for Precautionary Conservation Management of Wild Salmon. *Plos One* 8 (4): e60096. doi:10.1371/journal.pone.0060096
- Rooney RC & Podemski CL (2009) Effects of an experimental rainbow trout (*Oncorhynchus mykiss*) farm on invertebrate community composition. *Can J Fish Aquat Sci* 66(11): 1949-1964.
- Saksida SM (2006) Infectious haematopoietic necrosis epidemic (2001 to 2003) in farmed Atlantic salmon *Salmo salar* in British Columbia. *Dis Aquat Org* 72(3): 213-23.
- Saksida S, Constantine J, Karreman GA, Donald A (2007a) Evaluation of sea lice abundance levels on farmed Atlantic salmon (*Salmo salar* L.) located in the Broughton Archipelago of British Columbia from 2003 to 2005. *Aquacult Res* 38(3): 219-231.

# AQUACULTURE

## References

- Saksida S, Karreman GA, Constantine J, Donald A (2007b) Differences in *Lepeophtheirus salmonis* abundance levels on Atlantic salmon farms in the Broughton Archipelago, British Columbia, Canada. *J Fish Dis* 30(6): 357-366.
- Samuelsen OB, Lunestad BT, Husevag B, Holleland T, Ervik A (1992) Residues of oxolinic acid in wild fauna following medication in fish farms. *Dis Aquat Organ* 12: 111-119.
- Stechey D & Robertson WD (2010) An overview of closed-containment systems and technologies. In: Chadwick EMP, Parsons GJ, Sayavong B (eds) *Evaluation of closed-containment technologies for saltwater salmon aquaculture*. NRC Research Press, Ottawa, p 57-110.
- Tacon AGJ & Metian M (2008) Global overview on the use of fish meal and fish oil in industrially compounded aquafeeds: Trends and future prospects. *Aquaculture* 285(1-4): 146-158.
- Thorstad EB, Fleming IA, McGinnity P, Soto D, Wennevik V, Whoriskey F (2008) Incidence and impacts of escaped farmed Atlantic salmon *Salmo salar* in nature. NINA Special Report 36: 1-110. Accessed 27 Jul. <http://www.nina.no/archive/nina/PppBasePdf/temahefte/036.pdf>.
- Torrissen O, Jones S, Asche F, Guttormsen A, Skilbrei OT, Nilsen F, Horsberg TE, Jackson D (2013) Salmon lice - impact on wild salmonids and salmon aquaculture. *Journal of Fish Diseases* 36:171-194
- Tymchuk WE, Devlin RH, Withler RE (2006) The role of genotype and environment in phenotypic differentiation among wild and cultured salmonids. In: Fisheries and Oceans Canada (ed) *A Scientific Review of the Potential Environmental Effects of Aquaculture in Aquatic Ecosystems*. Volume IV. *Can Tech Rep Fish Aquat Sci* 2450: 4-10.
- Vike S, Nylund S, Nylund A (2009) ISA virus in Chile: evidence of vertical transmission. *Arch Virol* 154(1): 1-8.
- Volpe JP, Taylor EB, Rimmer DW, Glickman BW (2000) Evidence of natural reproduction of aquaculture-escaped Atlantic salmon in a coastal British Columbia river. *Conserv Biol* 14(3): 899-903.
- Waagbo R, Berntssen MHG, Danielsen T, Helberg H, Kleppa AL, Lea TB, Rosenlund G, Tvenning L, Susort S, Vikesa V, Breck O (2013) Feeding Atlantic salmon diets with plant ingredients during the seawater phase - a full-scale net production of marine protein with focus on biological performance, welfare, product quality and safety. *Aquaculture Nutrition* 19:598-618

# AQUACULTURE

## References

- Waddy SL, Burr ridge LE, Hamilton MN, Mercer SM, Aiken DE, Haya K (2002) Emamectin benzoate induces molting in American lobster, *Homarus americanus*. *Can J Fish Aquat Sci* 59: 1096-1099.
- Waddy SL, Merritt VA, Hamilton-Gibson MN, Aiken DE, Burr ridge LE (2007a) Relationship between dose of emamectin benzoate and molting response of ovigerous American lobsters (*Homarus americanus*). *Ecotox Environ Safe* 67(1): 95-99.
- Waddy SL, Mercer SM, Hamilton-Gibson MN, Aiken DE, Burr ridge LE (2007b) Feeding response of female American lobsters, *Homarus americanus*, to SLICE® -medicated salmon feed. *Aquaculture* 269(1-4): 123-129.
- White DG, McDermott PF (2009) Antimicrobial resistance in food borne pathogens. In: Jaykus LA, Wang HH, Schlesinger LS (eds) *Food Borne Microbes: Shaping the Host Ecosystem*. ASM Press, Washington, 231–265.
- Willis KJ & Ling N (2003) The toxicity of emamectin benzoate, an aquaculture pesticide, to planktonic marine copepods. *Aquaculture* 221: 289-297.